

STIC-ILL

370,193

NO

From: Flood, Michele  
Sent: Friday, November 02, 2001 6:07 PM  
To: STIC-ILL  
Subject: Following article needed. Thank you. CM1-11E11

09/646,740

TITLE: ISOFLAVONOIDS FROM IRIS-CROCEA.  
AUTHOR(S): SHAWL A S; KUMAR T  
SOURCE: PHYTOCHEMISTRY (OXF), (1992) 31 (4), 1399-1401.

TITLE: Determination of isoflavonoid glucosides in rhizomes of  
\*\*\*Belamcanda\*\*\* chinensis by high performance liquid  
chromatography.  
AUTHOR: Cui J.M.; Chung H.S.; Woo W.S.  
Korean J of Pharmacognosy (1993), 24/4 (309-312).

SOURCE: TITLE: Studies on chemical constituents of \*\*\*Belamcanda\*\*\*  
chinensis (L.)DC. II  
AUTHOR(S): Zhou, Li Xin; Lin, Mao  
SOURCE: Chin. Chem. Lett. (1997), 8(2), 133-134

TITLE: Pharmacological studies of \*\*\*tectoridin\*\*\* and  
\*\*\*tectorigenin\*\*\*  
AUTHOR(S): Esaki, Shunji  
SOURCE: Nippon Yakurigaku Zasshi (1968), 64(2), 186-98

BEST AVAILABLE COPY

Michele Flood  
Art Unit 1651  
CM 1-11E11  
308-9432

L.C.-R.C  
11/5/01-NAS

생약학회지  
Kor. J. Pharmacogn.  
24(4):309~312(1993)

## Determination of Isoflavonoid Glucosides in Rhizomes of *Belamcanda chinensis* by High Performance Liquid Chromatography

Jiong Mo Cui\*, Ha Sook Chung and Won Sick Woo

\*Department of Pharmacognosy, Yanbian Medical College, Yanji, China and  
Natural Products Research Institute, Seoul National University, Seoul 110-460, Korea

**Abstract**—A new method for separation and quantitative determination of isoflavone glucosides in rhizomes of *Belamcanda chinensis* (Iridaceae) by high performance liquid chromatography was elaborated. A reverse-phase system with a Spheri-5 RP-18 column using MeOH : HOAc : H<sub>2</sub>O (24 : 5 : 71) as a mobile phase was developed. The isoflavonoids were detected at 268 nm and the analysis was successfully carried out within 15 min.


**Keywords**—*Belamcanda chinensis* · Iridaceae · isoflavonoid glucosides · iridin · tectoridin · HPLC

The rhizomes of *Belamcanda chinensis* (Iridaceae) have been used as a Chinese drug for the treatment throat troubles such as tonsillitis<sup>(1)</sup>.

In the previous study, we examined the chemical constituents of this plant in order to isolate its active principles and to determine the structures of two major isoflavonoid glucosides such as iridin and tectoridin<sup>(2)</sup> together with a number of minor constituents<sup>(3-5)</sup>.

Recently, Wu and Xu reported an analytical method of tectoridin in *B. chinensis* by square wave voltammetry<sup>(6)</sup>. The present study describes a new method for separation of iridin and tectoridin and their simultaneous quantitative analysis by high performance liquid chromatography (HPLC).

According to our experiences, a reverse-phase column with a polar elution solvent mixture as a separation system has been established. It was revealed that major components in this plant, iridin and tectoridin (Fig. 1) could be separated



	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>
1	OH	OCH <sub>3</sub>	OCH <sub>3</sub>
2	H	OH	H

Fig. 1. Structure of iridin(1) and tectoridin(2)

with reasonable accuracy and this method may be employed for the estimation of flavone glucosides in pharmaceutical preparations of rhizomes of *B. chinensis*.

## Results and Discussion

### Selection of separation system

The rhizomes of *B. chinensis* were extracted

310




Fig. 2. HPLC chromatogram of the standard isoflavonoids mixture Iridin(I); Tectoridin(T). Analytical conditions; column, Spheri-5 RP-18, 5  $\mu$ m, 4.6 mm I.D.  $\times$  220 mm; eluent, MeOH : HOAc : H<sub>2</sub>O (24 : 5 : 71); flow rate, 1.0 ml/min; chart speed, 1.5 cm/min; detector, UV (268 nm). Concn.; 5.0  $\mu$ g/ml

with MeOH and the methanolic extract thus obtained was subjected to HPLC for iridin and tectoridin, monitoring with various solvent mixtures in order to find a good separation system. A reverse-phase system with a Spheri-5 RP-18 column using MeOH : HOAc : H<sub>2</sub>O (24 : 5 : 71) as a mobile phase was found to be applicable to the analysis of isoflavonoids in *B. chinensis*. The isoflavonoids were detected at 268 nm by UV detector.

Fig. 2 shows the HPLC chromatogram of a mixture of pure two isoflavonoids, which was obtained by elution with this solvent system. It can be seen that the baseline separation of the peaks obtained was relatively good.

Various parameters such as capacity factor of column, relative retention, resolution and retention time for the analytical conditions were calculated from the chromatogram (Fig. 2) and indicated in Table I.

Fig. 3 shows the typical HPLC chromato-

Kor. J. Pharmacogn.

Table I. HPLC parameters for isoflavonoids

Compounds	$t_R$	$t_R'$	$k'$	$\alpha$	R
Tectoridin	11.10	8.81	3.85	1.47	4.39
Iridin	14.06	11.77	5.65		

$t_R$ : retention time,  $t_R'$ : net retention time,  $k'$ : capacity factor of column,  $\alpha$ : relative retention, R: resolution.  $t_0$  = 2.29 min  
Solvent mixture; MeOH : HOAc : H<sub>2</sub>O (24 : 5 : 71), flow rate; 1.0 ml/min.




Fig. 3. HPLC chromatogram of methanol extract from domestic (A) and China (B) samples Iridin(I); Tectoridin(T)

The analytical conditions were the same as those in Fig. 2.

Concn.; A: 2.0  $\mu$ g/10 ml; B: 0.25  $\mu$ g/10 ml

grams of the methanolic extracts in the rhizomes of *B. chinensis*, domestics and imports from China. As shown in the chromatograms, the peaks appearing as high intensities are iridin (I) and tectoridin (T). It was evident that two compounds are major components of *B. chinensis* and solvent mixture and analytical conditions are very excellent in the chromatograms.

Vol. 24, No. 1, 1993

311




Fig. 4. Calibration curves for isoflavonoids

### Calibration curves

Based on chromatograms of varying amounts of pure isoflavonoids obtained by the reverse-phase system, their individual calibration curves were composed and shown in Fig. 4. The regression equation was calculated and statistical analysis was carried out to check for linearity of separation system.

Iridin:  $Y_1 = 90393X_1 - 8474$  (correlation coefficient,  $r_1 = 0.991$ );

Tectoridin:  $Y_2 = 90727X_2 + 7127$  (correlation coefficient,  $r_2 = 0.998$ ).

Each isoflavonoid component had a linear relationship between the amount ( $\mu\text{g} = X_1, X_2$ ) and peak area ( $Y_1, Y_2$ ). All the correlation coefficients of two calibration curves lay in the vicinity of the value of 1.00.

### Determination of isoflavonoids in the rhizomes of *B. chinensis*

A definite amount of the rhizomes of *B. chinensis* was pretreated by the method described in the experimental section and the content of isoflavonoids was analysed by the reverse-phase system. It was revealed that the contents of iridin and tectoridin in the rhizomes collected in Korea were analysed to be 0.3535% and 1.2140%, respectively, where as 0.4029% and 1.5350% in that imported from China. The method can be used to confirm isoflavonoids of *B. chinensis* not only quantitative but qualitative analysis.

## Experimental

**Plant material**—The rhizomes of *B. chinensis* were collected in the vicinity of Seoul. The Chinese origin was purchased from a drug store in Seoul and botanically identified.

**Apparatus**—The apparatus used was as follows: HPLC, Spectra Physics, Inc. attached with Spectra 100 variable wavelength detector; Injector, Rheodyne injection valve (10  $\mu\text{l}$ ) for HPLC; Column, Spheri-5 RP-18, dry-packed in 4.6 mm ID  $\times$  220 mm capillary column; Pump, SP 8800 ternary HPLC pump; Integrator, SP 4270 integrator.

**Reagents**—Reagents and extraction solvents were first grade.

Iridin, mp 208° and tectoridin, mp 257~258° were isolated from the rhizomes of *B. chinensis* according to the method of Lee, *et al.*<sup>2)</sup>

**Chromatographic conditions**—As a mobile phase, MeOH:HOAc:H<sub>2</sub>O (24:5:71) in reverse-phase system was used. The analysis was carried out at room temperature in flow rate of 1.0 ml/min. The substances eluted were detected by UV detector at wavelength of 268 nm. The chart speed was 1.5 cm/min.

**Preparation of calibration curves**—Each isoflavonoids (10 mg) was dissolved in 25 ml of MeOH. These solutions were further diluted stepwise with MeOH solution to 0.625, 1.25, 2.5 and 5.0  $\mu\text{g/ml}$ . And 10  $\mu\text{l}$  of each solution was injected on the column. The calibration curves were depicted by plotting the amounts of samples against peak areas.

**Analysis procedure**—Two grams of accurately weighed were extracted three times with 25 ml of boiling MeOH for 3 hr on a water bath and filtered. The methanolic solution was concentrated to dryness and lyophilized. The extract (10 mg) was dissolved in 25 ml of MeOH and diluted 0.25, 0.1 and 0.5  $\mu\text{g/10 ml}$ . And 10  $\mu\text{l}$

312

of each solution was applied and eluted to obtain the chromatogram.

(Received Sep. 16, 1993:

Accepted Oct. 8, 1993)

### References

1. Perry, L.M.: *Medicinal Plants of East and Southeast Asia*, MIT, Cambridge, p.181 (1980).
2. Lee, S.O., Woo, W.S., Woo, E.H. and Kim, K.S.: *Kor. J. Pharmacogn.* 20, 219 (1989).
3. Eu, G.H., Woo, W.S., Chung, H.S. and Woo, E.H.: *Kor. J. Pharmacogn.* 22, 13 (1991).
4. Chung, H.S. and Woo, W.S.: *Arch. Pharm. Res.* 14, 357 (1991).
5. Woo, W.S. and Woo, E.H.: *Phytochem.* 33, 939 (1993).
6. Wu, Y.X. and Xu, L.X.: *Acta Pharmaceutica Sinica* 27, 64 (1992).

*Kor. J. Pharmacogn.*

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**